


Combination thyroid hormone therapies treat hypothyroidism as well as levothyroxine

March 20 2021



Credit: CC0 Public Domain

Treatment of hypothyroidism, which results from an underactive thyroid gland, should be individualized and consideration should be given to alternatives to the first-line therapy, including desiccated thyroid extract



and combination therapy to replace the body's two main thyroid hormones, thyroxine (T4) and triiodothyronine (T3). Results of their new randomized clinical study are being presented at ENDO 2021, the Endocrine Society's annual meeting.

Combination therapy has been shown to be equally as effective as the standard treatment with levothyroxine alone, researchers say.

"There are now proven good treatment options for the more than one in 10 patients with hypothyroidism who continue to experience symptoms of fatigue, mental fogginess, weight gain and other symptoms despite taking levothyroxine," said the study's principal investigator, Thanh D. Hoang, D.O., a staff endocrinologist at Walter Reed National Military Medical Center in Bethesda, Md.

For some of these patients, Hoang said he has observed dramatic improvement—"a night and day difference"—in thyroid symptoms after they switch to desiccated thyroid extract (DTE), one of the drugs tested in the new study.

Also known as thyroid USP, DTE was the original treatment of hypothyroidism established almost 100 years ago. Unlike levothyroxine, which contains only one form of the thyroid hormone (thyroxine, T4), DTE comes from animal thyroids and contains both of the main thyroid hormones in the body, T4 and triiodothyronine (T3), Hoang said.

Past research from Hoang's group found that DTE safely led to modest weight loss and possible improvement in mental health over levothyroxine.

In their new study, Hoang and colleagues investigated the effectiveness of DTE in reducing thyroid symptoms as well as another combination therapy, levothyroxine plus liothyronine, the pharmaceutical grade



version of T4 and T3. Although a few studies have shown that this combination works better than levothyroxine monotherapy in some patients, other studies failed to show superiority. As a result, doctors do not always consider it when patients respond poorly to levothyroxine, Hoang said.

The investigators randomly assigned 75 adults with hypothyroidism to three months' treatment, with one of three drugs: DTE, levothyroxine-liothyronine combination, or levothyroxine alone. At the end of three months, the patients then randomly received a different treatment for three months, and finally took the third drug regimen for the last three months. Neither the patients nor the investigators were aware of which drug the patients were taking. After each treatment arm, patients completed a questionnaire of their thyroid symptoms.

The researchers found no significant difference between any of the three treatments in symptom relief. However, most patients reported a preference for combination therapy over levothyroxine alone, according to Hoang. Among the study participants, 45 percent reportedly preferred DTE over the other two treatments, 32 percent preferred the levothyroxine-liothyronine combination, and 23 percent preferred levothyroxine alone. Patients were more likely to prefer combination therapy if they had Hashimoto's disease, an autoimmune disease that is the most common cause of hypothyroidism in the U.S.

Because Hoang said it would be helpful for doctors to select which patients would initially respond better to a combination treatment than to levothyroxine alone, the researchers are further studying their data to find potential predictors of treatment response.

Provided by The Endocrine Society



Citation: Combination thyroid hormone therapies treat hypothyroidism as well as levothyroxine (2021, March 20) retrieved 20 September 2024 from https://medicalxpress.com/news/2021-03-combination-thyroid-hormone-therapies-hypothyroidism.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.