

C-section babies have microbiome deficit, but catch up over time

5 April 2021



intestine has already been colonized by bacteria and other microorganisms, and that the diversity of those species increases in the first few years of life.

This study provides more detailed information about development of the gut microbiome.

A key finding: Even at 5 years of age, it's incomplete.

How long it takes for the gut microbiome to mature differs from child to child, and this study found that babies delivered by C-section initially had less diverse gut bacteria than infants delivered vaginally.

But by the time they reached 3 to 5 years of age, the diversity of <u>gut bacteria</u> was similar in both groups, according to findings published online March 31 in the journal *Cell Host & Microbe*.

"It's striking that even at the age of 5 years, several of the bacteria that are important components of the intestinal <u>microbiota</u> in adults are missing in the children," Bäckhed said in a university news release.

That shows that the intestine is a complex and dynamic environment where <u>bacteria</u> create conditions for colonization by one another, he added.

"By investigating and understanding how the intestinal microbiota develops in healthy children, we may get a reference point to explore if the microbiota may contribute to disease in future studies," Bäckhed said.

More information: The Harvard School of Public Health has more on the <u>microbiome</u>.

Copyright © 2021 HealthDay. All rights reserved.

(HealthDay)—Infants born by cesarean section initially have less diverse gut bacteria than those delivered vaginally, but they catch up within a few years, new research reveals.

The researchers also found that it takes a long time for these bacteria colonies—known as the <u>gut</u> <u>microbiome</u>—to mature.

"Our findings show that the <u>gut microbiota</u> is a dynamic organ, and future studies will have to show whether the early differences can affect the cesarean children later in life," said study leader Fredrik Bäckhed. He's a professor of molecular medicine at the University of Gothenburg in Sweden.

For the study, his team took fecal samples from 471 newborns, and again when they were 4 months, 12 months, 3 years and 5 years old. The samples were analyzed for gut bacteria.

It was already known that at birth, an infant's



APA citation: C-section babies have microbiome deficit, but catch up over time (2021, April 5) retrieved 9 June 2021 from <u>https://medicalxpress.com/news/2021-04-c-section-babies-microbiome-deficit.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.