

New insights on cause of anaphylaxis following COVID-19 vaccination

7 April 2021



safe."

More information: P Sellaturay et al, Polyethylene glycol (PEG) is a cause of anaphylaxis to the Pfizer/BioNTech mRNA COVID?19 vaccine, *Clinical & Experimental Allergy* (2021). DOI: 10.1111/cea.13874

Provided by Wiley

A medical assistant prepares a dose of a COVID-19 vaccine to be administered to a patient. Credit: Public domain image courtesy of Lisa Ferdinando, U.S. Department of Defense

On the first day of the UK campaign for COVID-19 vaccination, there were reports of two cases of anaphylaxis—a severe, potentially life-threatening allergic reaction—within minutes of administration of the Pfizer/BioNTech mRNA vaccine. Subsequently further cases of suspected anaphylaxis to the Pfizer vaccine were reported.

A new report published in *Clinical & Experimental Allergy* reveals that an allergy to the ingredient polyethylene glycol (PEG) is a cause of anaphylaxis to the vaccine. However, this is in the <u>context</u> of millions of doses safely administered.

The authors note that very few people are allergic to PEG, and they provide a guide to identifying those who are at risk. "This preliminary report confirms PEG as a cause of anaphylaxis to the Pfizer/BioNTech vaccine, for the first time," they wrote. "COVID-19 vaccine anaphylaxis and PEG allergy are both rare, so proof of PEG as the cause in one case of vaccine anaphylaxis is important. However, it is important to emphasise that PEG allergy is rare and that COVID-19 vaccines remain



APA citation: New insights on cause of anaphylaxis following COVID-19 vaccination (2021, April 7) retrieved 6 November 2022 from https://medicalxpress.com/news/2021-04-insights-anaphylaxis-covid-vaccination.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.