

Exercise reduces risk of airway disease

April 27 2021



Credit: Pixabay/CC0 Public Domain

Exercise appears to reduce the long-term risk of bronchiectasis, a potentially serious disease of the airways, according to a study published in the journal *Radiology*.

Bronchiectasis is characterized by repeated cycles of inflammation and exacerbations that damage the airways, leaving them enlarged, scarred and less effective at clearing mucus. This creates an environment ripe for infections. Risk increases with age and the presence of underlying conditions like cystic fibrosis. There is no cure.

Computed tomography (CT) is used to confirm or rule out the [disease](#) in patients with symptoms like shortness of breath and coughing up mucus. Bronchiectasis has also been found on CT in asymptomatic and mildly symptomatic individuals.

Little is known about factors that can reduce the risk of bronchiectasis. While some studies have tied higher levels of [cardiorespiratory fitness](#) to a [reduced risk](#) of declining lung function and airway diseases, such as [chronic obstructive pulmonary disease](#), its benefits in reducing the risk of bronchiectasis are unknown.

To examine the association between [cardiorespiratory](#) fitness and bronchiectasis, researchers analyzed data from the long-running Coronary Artery Disease in Young Adults (CARDIA) study. CARDIA was launched in 1984 across the U.S. to examine the risk factors for [coronary artery disease](#) in [young adults](#).

The researchers looked at 2,177 healthy adults who were ages 18 to 30 years at the beginning of the study period. The study participants were followed up over a 30-year period with fitness tests and CT.

"We used year zero and year 20 cardiorespiratory fitness measured as exercise duration on a treadmill and ascertained bronchiectasis on chest CT at year 25," said study lead author Alejandro A. Diaz, M.D., M.P.H., assistant professor of medicine at Harvard Medical School and associate scientist at Division of Pulmonary and Critical Care Medicine at Brigham and Women's Hospital in Boston. "We assessed whether

differences in treadmill duration between year zero and 20 were associated with bronchiectasis on CT at year 25."

Of the 2,177 participants, 209, or 9.6%, had bronchiectasis at year 25. Preservation of cardiorespiratory fitness reduced the odds of bronchiectasis on CT at year 25.

"In an adjusted model, one minute longer treadmill duration between year zero and year 20 was associated with 12% lower odds of bronchiectasis on CT at year 25," Dr. Diaz said. "Having preserved fitness at middle age is associated with lower chances of bronchiectasis."

The researchers pointed to several possible explanations for the relationship between cardiorespiratory fitness and bronchiectasis. For one, a high level of cardiorespiratory fitness is linked with lower levels of systemic inflammation, which might help preserve the health of the airway. Good cardiorespiratory fitness also reduces the risk of certain diseases associated with bronchiectasis, such as asthma and pneumonia. Finally, high fitness levels may improve the ability of the [airway](#) to clear mucus.

The researchers observed a higher prevalence of bronchiectasis than found in previous studies. The difference may be explained by the use of CT for detecting bronchiectasis in the new study rather than the physician-based diagnosis used in previous studies.

"This study suggests that bronchiectasis on CT scans might be more frequent than previously thought," Dr. Diaz said. "However, the clinical implications of finding bronchiectasis on CT scans in people with no or mild symptoms remain to be determined."

The researchers are studying bronchiectasis in other populations like smokers to look for features of the airways and lung tissue associated

with bronchiectasis flare-ups.

"These results amplify the benefits of fitness to human health when a sedentary lifestyle is a concerning world epidemic," Dr. Diaz said. "It also highlights that fitness might be a tool to preserve lung health. The airways are challenged by what we breathe in every minute, and [fitness](#) may help to preserve lung health from injuries."

More information: Alejandro A. Diaz et al. Association between Cardiorespiratory Fitness and Bronchiectasis at CT: A Long-term Population-based Study of Healthy Young Adults Aged 18–30 Years in the CARDIA Study, *Radiology* (2021). [DOI: 10.1148/radiol.2021203874](#)

Provided by Radiological Society of North America

Citation: Exercise reduces risk of airway disease (2021, April 27) retrieved 6 April 2023 from <https://medicalxpress.com/news/2021-04-airway-disease.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|