

Guided yoga more effective than conventional treatment for preventing fainting

21 May 2021



Credit: CC0 Public Domain

Guided yoga is more effective than conventional treatment for preventing fainting in repeat sufferers, according to research published today in *EP Europace*, a journal of the European Society of Cardiology (ESC). Guided yoga therapy was also more beneficial in terms of quality of life than the standard treatment.

"Repeated fainting gives rise to mental stress, anxiety, depression, and a constant dread of the next episode," said study author Professor Jayaprakash Shenthar of the Sri Jayadeva Institute of Cardiovascular Sciences and Research, Bangalore, India. "Quality of life is negatively affected due to the unpredictability. Besides, some countries do not allow patients to drive."

Fainting or "[vasovagal syncope](#)" is a brief loss of consciousness caused by reduced blood flow to the brain. It can be triggered by prolonged standing, fear, pain, seeing blood, and hot, humid environments. It is estimated that approximately

half of the general population will have one syncopal event during their lifetime.

Recommendations to prevent vasovagal syncope include keeping hydrated, avoiding hot crowded environments, tensing the muscles, and lying down. "Unfortunately, most strategies have only modest benefits, which proves frustrating for patients," said Professor Shenthar.

This was the first randomized study to examine guided yoga as a treatment for patients with vasovagal syncope. The study enrolled 100 patients with at least three episodes of vasovagal syncope in the previous year. None of the patients were receiving treatment for syncope, and none were practicing yoga at the time of enrolment.

Participants were randomly allocated to the yoga or conventional treatment group. The conventional treatment group was advised to consume 6 to 9 grams of salt and more than 3 liters of water every day and perform physical counterpressure maneuvers such as crossing legs and tensing thigh and buttock muscles when warning signs of syncope occurred.

The yoga group was not given any specific advice regarding salt, fluid, or physical counterpressure maneuvers. Instead, they attended five to seven sessions in which a certified yoga instructor showed them how to perform a 60-minute yoga module. The yoga module consisted of pranayama (breathing), asanas (postures), and dhyana (meditation). Participants were then advised to perform yoga for 60 minutes at least five days per week for the subsequent 12 months.

Participants were asked to record the number of fainting episodes during follow-up. Quality of life was assessed at baseline, 3, 6, and 12 months

using the syncope functional status questionnaire—it asks how syncope affects [daily activities](#) such as driving, shopping, exercise, and relationships with family and friends and asks respondents to rate their fear/worry on a scale of 1 to 8.

The mean age of participants was 33 years, and more than half were women. Prior to the study, the average duration of syncopal episodes was 17 months, during which patients had an average of 6.4 blackouts. Compared to the conventional treatment arm, the yoga group had fewer blackouts and better quality of life scores at 3, 6, and 12 months. The [standard treatment](#) group reported 3.8 fainting episodes during 12 months of follow-up, while the yoga group reported 1.1 episodes.

Professor Shenthar said: "The benefits of guided yoga were apparent as early as three months after treatment initiation and were sustained at six and twelve months. Preliminary evidence suggests that the breathing and meditation aspects of yoga have a [positive effect](#) on the autonomic system while 'asanas' improve vascular tone—both of which could prevent gravitational pooling of blood in the lower parts of the body."

He concluded: "Our small, well-controlled, single-center study suggests that [yoga](#) may be a useful therapeutic option for patients with vasovagal syncope. Larger multicentre studies are needed to confirm the results."

More information: Jayaprakash Shenthar et al, A randomized study of yoga therapy for the prevention of recurrent reflex vasovagal syncope, *EP Europace* (2021). [DOI: 10.1093/europace/euab054](#)

Provided by European Society of Cardiology

APA citation: Guided yoga more effective than conventional treatment for preventing fainting (2021, May 21) retrieved 28 May 2021 from <https://medicalxpress.com/news/2021-05-yoga-effective-conventional-treatment-fainting.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.