

A fifth of asymptomatic COVID patients develop long COVID: study

June 15 2021



Credit: Pixabay/CC0 Public Domain

Almost a fifth of COVID patients without symptoms went on to experience conditions consistent with long COVID a month after their initial diagnosis, according to a huge study published Tuesday.

The analysis by non-profit FAIR Health encompassed [insurance claims](#) from 1.96 million Americans—the largest population of patients ever studied for long COVID—from February 2020 to February 2021.

"Even as the COVID-19 pandemic wanes, long-haul COVID persists as a public health issue affecting many Americans," said FAIR Health's president Robin Gelburd.

"The findings in our new study shed significant light on this emerging issue for all individuals who have long-haul COVID, as well as for [policy makers](#), providers, payors and researchers."

Perils of 'Long COVID'

"Long COVID" refers to symptoms of the disease that persist more than four weeks after being diagnosed.

The study found that across all ages, the most common post viral conditions were in order of frequency: pain, breathing difficulties, high cholesterol, general discomfort and fatigue, and [high blood pressure](#).

The odds of dying 30 days or more after initially being diagnosed with COVID were 46 times higher for patients who were hospitalized with COVID and discharged compared to those who weren't hospitalized.

Overall, 0.5 percent of COVID patients who were hospitalized then discharged died 30 days or more after their initial diagnosis.

Nineteen percent of asymptomatic COVID patients experienced long COVID symptoms 30 days out from their initial diagnosis; the figure grew to 27.5 percent of COVID patients who were symptomatic but not hospitalized, and 50 percent of those who were hospitalized.

The order of most common long COVID conditions varied by age group—for example in the pediatric population, intestinal issues replaced [high cholesterol](#) as the third most frequent.

Most long COVID conditions were associated more with females than males—but some, such as cardiac inflammation, were more common in males, who accounted for 52 percent of cases against 48 percent for females.

A quarter of all such cases occurred among individuals aged 19-29.

Among the four mental health conditions evaluated after 30 days, anxiety was the most common, followed by depression, adjustment disorders and tic disorder.

The biggest drawback of the new study is it lacks a control group of people who never got COVID, which would help determine the extent to which COVID caused the conditions as opposed to being coincidental.

The causes of long COVID, which is also known as long haul COVID, post-COVID syndrome or post-acute sequelae of COVID, remain unknown.

"Theories include persistent immune activation after the acute phase; initial damage from the virus, such as damage to nerve pathways, that is slow to heal; and persistent presence of low-level virus," the study said.

© 2021 AFP

Citation: A fifth of asymptomatic COVID patients develop long COVID: study (2021, June 15) retrieved 15 February 2024 from <https://medicalxpress.com/news/2021-06-asymptomatic-covid-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.