


Longer duration of positive COVID-19 PCR test results in people with certain comorbidities

16 June 2021



Novel Coronavirus SARS-CoV-2 Transmission electron micrograph of SARS-CoV-2 virus particles, isolated from a patient. Image captured and color-enhanced at the NIAID Integrated Research Facility (IRF) in Fort Detrick, Maryland. Credit: National Institute of Allergy and Infectious Diseases, NIH

New study results indicate that different comorbid conditions affecting individuals diagnosed with COVID-19 may impact how long they continue to receive positive SARS-CoV-2 polymerase chain reaction (PCR) test results. Individuals diagnosed with COVID-19 who are aged 60+, have three or more chronic medical conditions, particularly diabetes, obesity, rheumatologic disease, or an organ transplant, have positive PCR tests for longer periods of time compared to younger individuals without these comorbidities. However, the data showed no significant difference in the

duration of positive PCR tests results by the degree of immunocompromise or for individuals receiving chemotherapy or steroids to treat COVID-19.

Published online in *Open Forum Infectious Diseases* and led by researchers at Boston Medical Center, this study is the largest sample size to date of SARS-CoV-2 PCR retesting in both immunocompromised and non-immunocompromised patients, allowing for the comparison between these two groups.

"There are limited data on how long [immunocompromised individuals](#) continue to [test positive](#) for COVID-19 after initial infection," said Rachel Epstein, MD, MScE, an infectious diseases physician at Boston Medical Center and the study's corresponding author. "This study challenges whether the current recommendations to consider retesting severely immunocompromised individuals and use negative SARS-CoV-2 PCR test results to end isolation is necessary."

According to the Centers for Disease Control and Prevention (CDC), severely immunocompromised individuals are likely to have prolonged positive SARS-CoV-2 PCR test results. The conditions that cause immunocompromised status represented in the CDC definition include: people on chemotherapy or high dose steroids; those with untreated HIV; those with a primary immunodeficiency; or those who recently received an organ transplant.

The researchers examined data from 3,758 individuals who were retested using the SARS-CoV-2 PCR test following an initial positive result. The individuals were separated into groups by age as well as immunocompromised severity:

- Severe—active chemotherapy, HIV with a

CD4 count less than 200, organ transplant in removal from isolation necessary in severely immunocompromised individuals?, *Open Forum Infectious Diseases* (2021). DOI:

- Moderate—solid organ transplant recipient greater than 1 year prior, HIV with a CD4 count greater than 200, and others taking chronic biologic medications (4.2 percent of study group)
- Not immunocompromised (88.4 percent of study group)

Provided by Boston Medical Center

The median length of time for severely immunocompromised patients to receive a negative SARS-CoV-2 PCR test result was 22 days; for moderately and non-immunocompromised individuals, the time was 20 and 16 days, respectively. For individuals who had a solid [organ transplant](#), are older than 60, have diabetes, obesity or rheumatologic disease, as well as those with more than three comorbid conditions, it took longer for a negative SARS-CoV-2 PCR test result.

"Retesting individuals to end isolation precautions can delay care, and may not be necessary even for most severely immunocompromised people, particularly as most transmission studies demonstrate that it is highly unlikely for someone to transmit infection more than 20 days into their illness," said Epstein, an assistant professor of medicine and pediatrics at Boston University School of Medicine. The authors note that retesting recommendations should perhaps consider a combination of conditions or include only certain groups of extremely [immunocompromised](#) individuals. There have been a few cases in the literature showing positive COVID-19 test results months out from [initial infection](#) in patients who have undergone a solid organ or [bone marrow transplant](#) or received [chimeric antigen receptor \(CAR\) T-cell therapy](#).

Additional studies that directly measure transmission or transmissibility from individuals with a positive PCR test result more than 20 days into illness with COVID-19 would be helpful to better inform current CDC guidelines.

More information: Rachel L Epstein et al, Time to SARS-CoV-2 PCR clearance in immunocompromising conditions: is test-based

APA citation: Longer duration of positive COVID-19 PCR test results in people with certain comorbidities (2021, June 16) retrieved 3 September 2022 from <https://medicalxpress.com/news/2021-06-longer-duration-positive-covid-pcr.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.