


SARS-CoV-2 infections may trigger antibody responses against multiple virus proteins

21 June 2021



Credit: Pixabay/CC0 Public Domain

All coronaviruses produce four primary structural proteins and multiple nonstructural proteins. However, the majority of antibody-based SARS-CoV-2 research has focused on the spike and nucleocapsid proteins. A study published in *PLOS Biology* by Anna Heffron, Irene Ong and colleagues at the University of Wisconsin-Madison, U.S., suggests that immune responses may develop against other proteins produced by the SARS-CoV-2 virus.

The efficacy of spike [protein](#)-based vaccines is variable and not everyone infected with SARS-CoV-2 produces detectable antibodies against the spike or nucleocapsid proteins. Therefore, expanded antibody-based options have the potential to play an important role in improving vaccines, diagnostics, and therapeutics, particularly given the emergence of new variants. To investigate whether SARS-CoV-2 infection induces robust antibody responses against all SARS-CoV-2 proteins, researchers mapped 79 "epitopes"—specific regions of the viral proteome that antibodies recognize and bind to. They also tested whether antibodies that develop in response to SARS-CoV-2 or existing antibodies from

previous exposures to coronaviruses might bind to any of the proteins in the six other known human coronaviruses to identify potential cross-reactive epitopes.

In addition to spike and nucleocapsid proteins, the authors located previously unknown, highly reactive B cell epitopes throughout the full array of proteins in SARS-CoV-2 and other coronaviruses, expanding the potential for future [vaccine](#) and therapeutic development. Future research is needed, however, to determine how long these [antibodies](#) remain and whether responses of vaccinated individuals differ from those who contracted COVID-19 prior to vaccination. Dr. Ong and colleagues will continue to investigate these aspects in adults and children.

Although the authors did not directly profile variants of concern that have emerged since the beginning of the COVID-19 pandemic, a comparison of the original SARS-CoV-2 genome with a few of the variants of concern identified numerous variations in regions that are at or within 3 [amino acids](#) of identified antibody binding epitopes.

According to the authors, "Our extensive profiling of epitope-level resolution antibody reactivity in COVID-19 convalescent subjects, confirmed by independent assays, provides new epitopes that could serve as important targets in the development of improved diagnostics, vaccines, and therapeutics against SARS-CoV-2, variants of concern, and dangerous human coronaviruses that may emerge in the future."

More information: Anna S. Heffron et al, The landscape of antibody binding in SARS-CoV-2 infection, *PLOS Biology* (2021). [DOI: 10.1371/journal.pbio.3001265](https://doi.org/10.1371/journal.pbio.3001265)

Provided by Public Library of Science

APA citation: SARS-CoV-2 infections may trigger antibody responses against multiple virus proteins (2021, June 21) retrieved 12 October 2022 from <https://medicalxpress.com/news/2021-06-sars-cov-infections-trigger-antibody-responses.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.