


# Distinctive gut microbiome unrelated to diet may characterize children with autism

July 26 2021

---



Credit: CC0 Public Domain

Children with autism seem to have a distinctive and underdeveloped range and volume of gut bacteria (microbiome) that isn't related to their diet, suggests a small study published online in the journal *Gut*.

They have significantly fewer [bacteria](#) linked to neurotransmitter activity and 5 [species of bacteria](#) that aren't typically found in the guts of [children](#) without the condition, suggesting that there may be a characteristic microbial profile for [autism](#), which may pave the way for treatment early on, say the researchers.

Apart from [genetic factors](#), it has been suggested that the [gut microbiome](#) may have a part to play in autism spectrum disorders. And the evidence suggests that the pathway between gut bacteria and the central nervous system, referred to as the gut-brain axis, has a profound effect on social behaviours.

Diagnosing autism spectrum disorder can be difficult, because there is no definitive medical test and diagnosis is based on physician assessment.

The researchers therefore wanted to see if 3-6 year olds with autism might harbour a [microbiome](#) that differs significantly from that of typically developing children, which might be used to facilitate early treatment.

They compared the range, volume, and associated functions of bacteria in the stool samples of 128 Chinese children, 64 of whom had autism spectrum disorder and 64 of whom didn't.

The factors most strongly and independently associated with microbiome composition in the stool samples were age, autism, and weight (BMI). Diet wasn't an influential factor.

The stool samples of children with autism displayed a greater variety of microbes than those matched for age and weight who didn't have the condition.

Clostridium, Dialister and Coprobacillus were enriched in children with autism while Faecalibacterium was significantly decreased.

Several Clostridium species enriched in children with autism closely interacted with each other and formed a connected group. Clostridia species have been linked with autism via the production of clostridial toxins which can damage the central nervous system, point out the researchers.

And the gut microbiome composition in children with autism differed substantially from that of children without the condition, with five [bacterial species](#) largely accounting for the difference: Alistipes indistinctus; candidate division\_TM7\_isolate\_TM7c (single cell organism); Streptococcus cristatus; Eubacterium limosum; and Streptococcus oligofermentans.

These differences were further confirmed in a separate group of 18 children, 8 with autism and 10 without.

Importantly, bacteria associated with neurotransmitter activities were substantially reduced in children with autism.

Some 26 age-related bacterial species were identified as proxies of typical development of the gut microbiome by age. But these associations were absent in children with autism, suggesting abnormal

development of the gut microbiome during the early life of these children, say the researchers.

This is a small study and the gut microbiome can vary according to geographical region. Nevertheless, the researchers write: "Our study shows for the first time that the gut microbiota of children with [autism] is abnormally developed and lags that of age-matched peers.

"As development of microbial communities within the [gastrointestinal] tract during childhood represents a critical window of human growth and health, shifts in the gut microbiota during early life development may have important functional roles in the pathogenesis of [autism spectrum disorder] and thus warrants extensive investigation."

And they conclude: "Our study supports the potential role of non-invasive prediction of [autism spectrum disorder] based on faecal bacteria markers and age-related bacteria development profile.

"Future therapeutics targeting reconstitution of gut microbiota in early life and increasing abundance of neurotransmitter-synthesised bacteria such as *Faecalibacterium* should be explored for [[autism spectrum disorder](#)]."

**More information:** Underdevelopment of the gut microbiota and bacteria species as non-invasive markers of prediction in children with autism spectrum disorder, *Gut* (2021). [DOI: 10.1136/gutjnl-2020-324015](#)

Provided by British Medical Journal

Citation: Distinctive gut microbiome unrelated to diet may characterize children with autism

(2021, July 26) retrieved 30 December 2023 from

<https://medicalxpress.com/news/2021-07-distinctive-gut-microbiome-unrelated-diet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.