

Study provides new insights on tree nut allergies

18 August 2021



Prevalence and early-life risk factors for tree nut sensitization and allergy in young adults, *Clinical & Experimental Allergy* (2021). DOI: [10.1111/cea.13994](https://doi.org/10.1111/cea.13994)

Provided by Wiley

Credit: CC0 Public Domain

In a study published in *Clinical and Experimental Allergy*, allergic sensitivity to tree nuts was common in Swedish adults, but most people never experienced symptoms.

In the study of 2,215 individuals, eczema, asthma, and [egg allergies](#) at an early age were associated with developing a tree nut allergy by adulthood. Also, allergen molecules were better [diagnostic tools](#) for predicting allergic symptoms to tree nuts compared with analyzing allergen extracts.

Tree nuts include hazelnuts, walnuts, pecans, cashews, pistachios, almonds, and brazil nuts.

"This study increases the understanding of tree nut allergy in a [general population](#), followed from infancy up to adulthood. For example, our study reveals that most extract-based tree nut-sensitized individuals do not have tree nut allergy and hence extract-based testing for tree nuts without a specific clinical suspicion should not be performed," said co-lead author Jessica Bager, of the Karolinska Institutet, in Sweden.

More information: Jessica Bager et al,

APA citation: Study provides new insights on tree nut allergies (2021, August 18) retrieved 1 July 2022 from <https://medicalxpress.com/news/2021-08-insights-tree-nut-allergies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.