

# Human health may be at risk from long-term exposure to air pollution below current air quality standards and guidelines

September 1 2021

---



Credit: CC0 Public Domain

Long-term exposure to air pollution appears to still be linked to higher mortality despite the existence of air quality standards that restrict levels

of pollution, suggests a study published online in *The BMJ* today.

Researchers found evidence of higher [death](#) rates amongst people who had been exposed to more air [pollution](#) even though the levels were allowed under current official standards.

Previous studies have found an association between long term exposure to outdoor air pollution such as those in the form of fine particles in the air (known as [particulate matter](#) or PM<sub>2.5</sub>) and [nitrogen dioxide](#) (NO<sub>2</sub>) and poor health or death.

Air pollution concentrations have fallen substantially in Europe since the 1990s, but it is unclear whether there still is a link between pollution and ill health or death at concentrations of pollution that are below current permitted limits.

Therefore, an international team of researchers led by the Institute for Risk Assessment Sciences at Utrecht University in the Netherlands, set out to investigate if there was an association between low levels of air pollution concentrations and natural and cause specific deaths.

Low level air pollution was defined as concentrations below current limit values as set by the European Union, US Environmental Protection Agency and the World Health Organization (WHO) air quality guidelines.

The researchers analysed data on eight groups of people within six European countries—Sweden, Denmark, France, the Netherlands, Germany and Austria—totalling 325,367 adults collectively.

Their study, known as the Effects of Low-Level Air Pollution: A Study in Europe (ELAPSE) recruited participants in the 1990s or 2000s.

Of the 325,367 participants who were followed up over an almost 20-year period, around 14.5% (47,131 people) died during the study period.

Analysis of the results showed that people who had higher exposure to particulate matter (PM<sub>2.5</sub>), nitrogen dioxide, and black carbon were more likely to die.

An increase of 5 µg/m<sup>3</sup> (a concentration measure of particulate matter) in PM<sub>2.5</sub> was associated with a 13% increase in natural deaths while the corresponding figure for a 10 µg/m<sup>3</sup> increase in nitrogen dioxide was 8.6%. Associations with PM<sub>2.5</sub> and nitrogen dioxide were largely independent of each other.

Moreover, associations with PM<sub>2.5</sub>, nitrogen dioxide, and black carbon remained significant at low to very low concentrations.

For people who were exposed to pollution levels below the US standard of 12 µg/m<sup>3</sup>, an increase of 5 µg/m<sup>3</sup> in PM<sub>2.5</sub> was associated with a 29.6% increase in natural deaths. People exposed to nitrogen dioxide at less than half the current EU standard of 40 µg/m<sup>3</sup>, a 10 µg/m<sup>3</sup> increase in [nitrogen dioxide](#) was associated with a 9.9% increase in natural deaths.

This is an observational study, and as such, can't establish cause.

The study also has some limitations, say the authors, such as the fact that it focused on exposure in 2010 which was towards the end of the follow-up period for most participants and, given the downward trend in air pollution, this measure might not exactly reflect the concentrations experienced during follow-up.

However, this was a large study from multiple European groups of

people with detailed information provided.

As such, the authors conclude: "Our study contributes to the evidence that outdoor [air pollution](#) is associated with mortality even at levels below the current European and North American standards and WHO guideline values.

"These findings are therefore an important contribution to the debate about revision of air quality limits, guidelines and standards, and future assessments by the Global Burden of Disease [study]."

**More information:** Long term exposure to low level air pollution and mortality in eight European cohorts within the ELAPSE project: pooled analysis, *BMJ* (2021). [DOI: 10.1136/bmj.n1904](https://doi.org/10.1136/bmj.n1904)

Provided by British Medical Journal

Citation: Human health may be at risk from long-term exposure to air pollution below current air quality standards and guidelines (2021, September 1) retrieved 6 May 2023 from <https://medicalxpress.com/news/2021-09-human-health-long-term-exposure-air.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--