


What are ideal blood sugar levels for preventing repeat strokes, heart attacks?

September 29 2021



Credit: CC0 Public Domain

Blood sugar control has always been important for people with diabetes when it comes to preventing a stroke. But a new study finds for people

with diabetes who have a stroke, there may be an ideal target blood sugar range to lower the risk of different types of vascular diseases like a stroke or heart attack later on. The research is published in the September 29, 2021, online issue of *Neurology*, the medical journal of the American Academy of Neurology.

"We know that having diabetes may be associated with an increased risk of having a first [stroke](#)," said study author Moon-Ku Han, MD, Ph.D., of Seoul National University College of Medicine in Korea. "But our results indicate that there is an optimal blood sugar level that may start to minimize the risk of having another stroke, a heart attack or other [vascular problems](#), and it's right in the 6.8% to 7.0% range."

The study involved 18,567 people with diabetes with an average age of 70. All participants were admitted to the hospital for an [ischemic stroke](#), which is caused by a blood clot. Upon admission, researchers used a test called the hemoglobin A1C to determine people's average blood sugar level over the past two to three months. This test measures a percentage of hemoglobin proteins in the blood coated with sugar. A level below 5.7% is considered normal; 6.5% or higher generally indicates diabetes. The participants had an average A1C of 7.5%.

Researchers then followed up one year later to find out if there was an association between A1C levels with the risk of having another stroke, a heart attack, or dying from these or other vascular causes.

Of all participants, 1,437, or about 8%, had a heart attack or died from vascular disease within a year of starting the study, and 954, or 5%, had another stroke.

The study found that people admitted to the hospital with A1C levels above the 6.8% to 7.0% range had an [increased risk](#) of having a vascular event like a heart attack, as well as having another stroke.

After adjusting for factors like age and sex, researchers found that people's risk for a [heart attack](#) or similar vascular diseases was 27% greater when they were admitted to the hospital with A1C levels above 7.0%, compared to those admitted with A1C levels below 6.5%. People's risk for having another stroke was 28% greater when admitted to the hospital with A1C levels above 7.0%, compared to those below 6.5%.

"Our findings highlight the importance of keeping a close eye on your blood sugar if you're diabetic and have had a stroke," Han said.

A limitation of the study is that people's blood sugar levels were measured only at the start of the study; no follow-up levels were available.

Provided by American Academy of Neurology

Citation: What are ideal blood sugar levels for preventing repeat strokes, heart attacks? (2021, September 29) retrieved 15 January 2024 from <https://medicalxpress.com/news/2021-09-ideal-blood-sugar-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--