




Patients with severe COVID-19 could benefit from higher doses of corticosteroids

October 22 2021

A No. of days alive without life support at 28 d as horizontally stacked proportions



B Time to death curves censored at 90 d



No. at risk		0	10	20	30	40	50	60	70	80	90
12 mg of dexamethasone		497	430	384	357	344	342	337	334	333	333
6 mg of dexamethasone		485	416	365	322	314	305	301	300	299	298

Distributions of the Primary Outcome and Time to Death Curves to Day 90 A, Life support was defined as invasive mechanical ventilation, circulatory support, or kidney replacement therapy. There were missing data in 11 patients for the

primary outcome. Red represents the worse outcomes and blue represents better outcomes. B, There were 14 patients who were not followed up for the full 90 days (7 patients in each intervention group) and who were included until the last day they were known to be alive. The median follow-up time was 90 days (IQR, 24-90 days) in the 12 mg of dexamethasone group and 90 days (IQR, 20-90 days) in the 6 mg of dexamethasone group. The time to death was compared post hoc using unadjusted Cox regression. Credit: DOI: 10.1001/jama.2021.18295

A large international study in hospitalized patients with severe COVID-19 has shown that while a higher dose of steroids did not significantly reduce mortality, there was a trend towards benefit without increased side effects.

Published today in the *Journal of the American Medical Association*, the study compared the standard 6mg dose of the steroid dexamethasone with 12mg in patients requiring a higher degree of support to maintain their oxygen levels. It is the first COVID-19 trial to report on long term mortality and one of only a few that were "blinded," or designed in a way so that doctors didn't know which patient received which dose.

Study co-author Professor Bala Venkatesh from The George Institute for Global Health said that although previous research suggested higher doses of dexamethasone may benefit patients with more severe COVID-19, there had been concerns about potential adverse reactions.

"Our study found no differences in the adverse event rates between the two doses. While there had been reports of infections with so called 'black fungus' in patients with COVID-19 with a weakened immune system, our study found no increase in infection rates with the higher doses of steroids," he said.

Patients with critical COVID-19 typically have severe lung inflammation

and very [low oxygen levels](#), which often leads to the requirement for increasing oxygen support, mechanical ventilation, support to maintain blood pressure and kidney dialysis. Dexamethasone is the mainstay of treatment for severe COVID-19. While a 6mg daily dose is recommended for up to ten days, there have been indications that a higher dose may benefit those with more severe disease. A potential side-effect of steroid use is suppression of the body's [immune system](#), diminishing the ability to fight other types of infections. There have been some reports of steroid-treated COVID-19 patients developing serious fungal infections, known as mucormycosis or "black fungus."

In this study, researchers recruited 1000 adult patients with confirmed SARS-CoV-2 [infection](#) across 31 sites in 26 hospitals in Denmark, India, Sweden, and Switzerland between August 2020 and May 2021. The patients were randomly assigned into two groups, one receiving intravenous dexamethasone 12 mg (n=503) and the other dexamethasone 6 mg daily (n=497) for up to ten days. The proportion of patients alive and not requiring life support after 28 days were 42.6 percent in the 12mg group and 40.2 percent in the 6mg group. The [death rates](#) were 27.1 percent and 32.3 percent in patients assigned to 12mg and 6mg group, respectively.

The George Institute India's Professor Vivek Jha, who along with Prof Venkatesh set up and conducted the trial in the 12 participating hospitals in India, said that the study provided useful guidance to clinicians treating COVID-19 patients around the world.

"Whilst the data do not provide unequivocal evidence that dexamethasone 12mg is better than 6 mg, we saw a trend towards reduced requirement for life support and mortality at the higher dose without any increase in risk of serious infections," he said. "As [dexamethasone](#) is cheap, easily available and indicated for the treatment of COVID-19 [patients](#) with critically low [oxygen levels](#), even a small

difference in death rates or health outcomes could lead to important clinical and health economic benefits at the population level."

More information: Effect of 12 mg vs 6 mg of Dexamethasone on the Number of Days Alive Without Life Support in Adults With COVID-19 and Severe Hypoxemia, *JAMA* (2021). [DOI: 10.1001/jama.2021.18295](https://doi.org/10.1001/jama.2021.18295)

Provided by George Institute for Global Health

Citation: Patients with severe COVID-19 could benefit from higher doses of corticosteroids (2021, October 22) retrieved 14 April 2023 from <https://medicalxpress.com/news/2021-10-patients-severe-covid-benefit-higher.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.