

Stress, by itself, can lead to excessive drinking in women but not men

December 13 2021



Credit: CC0 Public Domain

A new study has shown that stress alone can drive women to excessive drinking.

Men who experienced the same stress only drank to excess when they had already started consuming [alcohol](#).

Though rates of alcohol misuse are [higher in men than women](#), [women](#) are [catching up](#). Women also have [a greater risk](#) than men of developing alcohol-related problems.

Participants consumed alcoholic beverages in a simulated bar while experiencing stressful and non-stressful situations. Stress led women, but not men, to drink more than intended, a finding that demonstrates the importance of studying sex differences in [alcohol consumption](#). The study was published in *Psychology of Addictive Behaviors*.

"Some people can intend to have one or two [alcoholic beverages](#) and stop drinking, but other people just keep going. This impaired control over drinking is one of the earliest indicators of alcohol use disorders, and we know stress contributes to both impaired control over drinking and dysregulated drinking. The role of stress in impaired control over drinking is understudied, especially in women," said Julie Patock-Peckham, assistant research professor at ASU and lead author on the study.

The study took place in a research laboratory designed to simulate a bar, complete with a bartender, bar stools and lively conversations. The participants included 105 women and 105 men. They were randomized into different groups, with some either experiencing a stressful situation and others a non-stressful situation. Next, half the participants received an alcoholic drink that was equivalent to three cocktails, and the other half received three non-alcoholic drinks. After that, all participants had unrestricted access to [alcoholic drinks](#) from the bar for 90 minutes.

"We know that both genes and the environment play a role in problematic drinking. We can't do anything about the genes, but we can

intervene with the environment. Stress and impaired control over drinking are tightly connected, and because stress is something we can manipulate, we tested whether stressors cause dysregulated drinking," said Patock-Peckham, who leads the Social Addictions Impulse Lab at ASU.

The experimental set-up let the research team determine whether stress, the initial drink or the combination of the two caused how much alcohol the participants consumed. The team measured alcohol consumption in total number of drinks consumed and by using breath blood alcohol content (BAC).

Exposure to stress led to heavier drinking in all participants. Men who received a first drink with alcohol in it and experienced stress drank more than men who received the placebo.

Whether the first drink was alcoholic or not did not matter for women: Experiencing stress led to heavy [drinking](#).

"That women just needed the [stress](#) but men needed the push of already having alcohol on board shows how important this type of research is," Patock-Peckham said. "The outcomes from alcohol use are not the same for men and women, and we cannot keep using models that were developed in men to help women."

More information: Effects of stress, alcohol prime dose, and sex on ad libitum drinking, *Psychology of Addictive Behaviors* (2021). [DOI: 10.1037/adb0000801](https://doi.org/10.1037/adb0000801)

Provided by Arizona State University

Citation: Stress, by itself, can lead to excessive drinking in women but not men (2021, December 13) retrieved 24 February 2023 from <https://medicalxpress.com/news/2021-12-stress-excessive-women-men.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.