

Response to exercise is key to novel device therapy for the most common type of heart failure

February 1 2022



Credit: Pixabay/CC0 Public Domain

Heart failure with preserved ejection fraction (HFpEF), also called diastolic heart failure, affects 3 million Americans. Despite being the most common type of heart failure in the United States, effective

treatments remain elusive, leading to high morbidity and mortality.

"HFpEF makes up half all [heart failure](#) cases, yet we have very limited treatment options," said Sanjiv Shah, MD, director of research at the Bluhm Cardiovascular Institute and director of the Northwestern Medicine HFpEF Program. "Most standard therapies for [heart](#) failure are ineffective in this condition, leaving a major unmet need for a large patient population."

This type of heart failure occurs when the [left ventricle](#) is unable to relax, limiting the amount of blood filling into the heart, which causes fluid to build up in the lungs and the body, causing symptoms including shortness of breath, fluid retention, irregular heartbeat, and exercise intolerance.

A Northwestern Medicine-led study published in *The Lancet* suggests that some patients with HFpEF may benefit from a novel, minimally invasive cardiac implant device called an atrial shunt. The study also offers new insight into the role exercise plays in understanding, diagnosing and treating this type of heart failure.

"While the overall trial was neutral, in our subgroup analyses, we found that what happens in the heart and lungs during exercise is of prime importance in this type of heart failure," said Dr. Shah, international principal investigator of the trial. "The normal response to exercise is relaxation of the blood vessels in the lungs. Patients with HFpEF who are able to relax the blood vessels in their lungs appear to do well with the device, whereas those whose blood vessels can't relax appear to do worse when an atrial shunt is implanted."

An atrial shunt is placed through a catheter, creating a small hole between the left and right atria allowing blood to flow from the stiff left atrium to the normal right atrium, potentially lowering pressure in the

left atrium and reducing the symptoms of HFpEF. The procedure is minimally invasive, low-risk and patients go home the next day.

"What we saw in this study is encouraging and suggests that future clinical [trials](#) should specifically investigate the subgroup of patients with HFpEF whose pulmonary blood vessels respond normally to exercise," said Dr. Shah. "If future trials validate what we found, the potential is enormous. This subgroup comprises two thirds of people with this type of heart failure—that is 2 million people could benefit from this innovative therapy. This simple, one-time procedure could reduce hospitalizations and significantly improve quality of life."

While cardiovascular conditions such as coronary artery disease are routinely diagnosed with exercise testing, clinical assessment for HFpEF is done at rest—something that Dr. Shah hopes will change following this trial.

"This has potential to change the way we evaluate patients with this condition and should guide how future clinical trials are conducted and the criteria for enrollment," said Dr. Shah.

The Corvia REDUCE LAP-HF II pivotal trial is the largest device study ever conducted in HFpEF and the first pivotal trial of interatrial shunts completed.

More information: Sanjiv J Shah et al, Atrial shunt device for heart failure with preserved and mildly reduced ejection fraction (REDUCE LAP-HF II): a randomised, multicentre, blinded, sham-controlled trial, *The Lancet* (2022). [DOI: 10.1016/S0140-6736\(22\)00016-2](https://doi.org/10.1016/S0140-6736(22)00016-2)

Provided by Northwestern Memorial HealthCare

Citation: Response to exercise is key to novel device therapy for the most common type of heart failure (2022, February 1) retrieved 4 February 2024 from <https://medicalxpress.com/news/2022-02-response-key-device-therapy-common.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.