


COVID-19 clinical trials disproportionately run out of high-income countries

February 28 2022



Ratio of participant recruitment to COVID-19 cases by country and income status in relation to total COVID-19 cases. Bubbleplot showing participant recruitment to COVID-19 cases ratio stratified by income status and COVID-19



burden. Each bubble represents one country, with countries color-coded by income status. The size of each bubble is proportional to the ratio of participant recruitment to COVID-19 cases in that country. A larger bubble therefore means that that country has recruited a larger number of participants into trials relative to the total number of COVID-19 cases in that country. Overall, there were 80 high-income countries, 56 upper-middle-income countries, 55 lower-middle countries, and 27 low-income countries, of which 23 high-income, 8 upper-middle-income, 3 lower-middle-income, and 0 low-income countries recruited patients. The remaining countries with no participants recruited in COVID-19 trials are represented as small black circles. Credit: DOI: 10.1001/jamanetworkopen.2022.0444

New research has shown that COVID-19 clinical trials are being disproportionately run out of high-income countries.

Published in *Jama Network Open* and led by researchers from the Doherty Institute and The George Institute, the study found the majority of randomized clinical <u>trials</u> for COVID-19 are being performed in high-income countries, clearly misrepresenting the global burden of the disease.

Professor Steven Tong, a Royal Melbourne Hospital infectious diseases clinician, co-lead of clinical research at the Doherty Institute, Principal Investigator of the AustralaSian COVID-19 Trial (ASCOT) and senior author on the paper, said the inequality could bring into doubt the validity and generalisability of these trials results.

"A treatment that works in an <u>intensive care unit</u> in the United States is not necessarily going to work in low and middle income countries like India, where substantial differences in healthcare capacity and other population factors exist," said Professor Tong.



"It brings into question how much these results can be applied to low and middle income countries."

Of the 295,845 participants who were enrolled in a randomized clinical trial, 84 percent were recruited from <u>high-income countries</u>.

What's more, of the 16 percent who were not from high income countries, none were from low income countries, with these participants either from lower middle or upper middle income countries.

This is in stark contrast to where the burden of disease for COVID-19 fell, with 58% of reported cases from low and middle income countries up to mid 2021.

Professor Bala Venkatesh from The George Institute and on the international trial steering committee of ASCOT, said that despite a huge COVID-19 disease burden and resource limitations, India has made a major contribution to COVID-19 research.

"COVID-19 has amplified the inequalities in global health and this clearly extends to who leads and who participates in randomized <u>clinical</u> <u>trials</u>," Professor Tong said.

"We need to now look at prioritizing research that is relevant to low and middle income countries, and building research infrastructure to support the conduct of such trials."

More information: Mahesh Ramanan et al, Geographical Representation of Low- and Middle-Income Countries in Randomized Clinical Trials for COVID-19, *JAMA Network Open* (2022). DOI: 10.1001/jamanetworkopen.2022.0444



Provided by George Institute for Global Health

Citation: COVID-19 clinical trials disproportionately run out of high-income countries (2022, February 28) retrieved 12 February 2024 from https://medicalxpress.com/news/2022-02-covid-clinical-trials-disproportionately-high-income.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.