

Eating protein from a greater variety of sources may lower risk of high blood pressure

March 10 2022



Credit: Unsplash/CC0 Public Domain

Eating a balanced diet including protein from a greater variety of sources may help adults lower the risk of developing high blood pressure,

according to new research published today in *Hypertension*, a peer-reviewed journal of the American Heart Association.

Nearly half of the U.S. population has [hypertension](#), or [high blood pressure](#)—one of the leading contributors to cardiovascular disease. When left untreated, high blood pressure damages the circulatory system and is a significant contributing factor to [heart attack](#), stroke and other [health conditions](#).

"Nutrition may be an easily accessible and effective measure to fight against hypertension. Along with fat and carbohydrates, protein is one of the three basic macronutrients," said study author Xianhui Qin, M.D., of the National Clinical Research Center for Kidney Disease at Nanfang Hospital, Southern Medical University in Guangzhou, China.

There is a strong association between poor diet quality and increased risk of cardiovascular disease and death from cardiovascular disease. In its [2021 dietary guidance](#) to improve cardiovascular health, the American Heart Association advises people eat healthy sources of protein, mostly from plants and may include seafood and low-fat or fat-free dairy products, and, if desired, lean cuts and unprocessed forms of meat or poultry. The American Heart Association recommends eating one to two servings, or 5.5 ounces, of protein daily.

The study authors analyzed health information for nearly 12,200 adults living in China who were part of at least 2 out of 7 rounds of the China Health and Nutrition Survey from 1997 to 2015 (surveys taken every 2-4 years). Participants' initial survey was used as a baseline, while data from their last round was used as a follow-up for comparison. Participants were an average age of 41 years, and 47% were men. The survey measured [dietary intake](#) in three consecutive 24-hour dietary recalls and a household food inventory. A trained interviewer collected 24-hour dietary information over 3 days in the same week during each round of

the survey.

Participants were given a protein "variety score" based on the number of different sources of protein eaten out of 8 reported: whole grains, refined grains, processed red meat, unprocessed red meat, poultry, fish, egg and legumes. One point was given for each source of protein, with a maximum variety score of 8. The researchers then evaluated the association for new onset hypertension in relation to the protein variety score.

New-onset hypertension was defined as systolic (top number) blood pressure greater than or equal to 140 mm Hg and/or diastolic (bottom number) blood pressure greater than or equal to 90 mm Hg, taking blood pressure-lowering medicine, or self-reporting that a physician diagnosed high blood pressure since their last survey visit. Average time to follow-up was 6 years.

The analysis found:

- More than 35% of the nearly 12,200 participants developed new-onset high hypertension during follow-up.
- Compared to participants with the lowest variety score for protein intake (less than 2), those with the highest variety score (4 or higher) had a 66% lower risk of developing high blood pressure.
- For each of the 8 protein types, there was a window of consumption amount where the risk of hypertension was lower. Researchers described this as the appropriate level of consumption.
- When total quantity of protein intake was considered, the amount consumed was divided into five categories (quintiles), from least to most intake. People who ate the least amount of total protein and those who ate most protein had the highest risk for new onset

of hypertension.

"The heart health message is that consuming a [balanced diet](#) with proteins from various different sources, rather than focusing on a single source of dietary protein, may help to prevent the development of high blood pressure," Qin said.

A limitation of the study is its observational design. Because researchers used prior [health information](#), they could not definitively prove [protein](#) intake of any kind or quantity caused or prevented new-onset hypertension.

Co-authors are Chun Zhou, M.S.; Qimeng Wu, M.S.; Ziliang Ye, M.S.; Mengyi Liu, M.S.; Zhuxian Zhang, M.S.; Yuanyuan Zhang, M.D.; Huan Li, M.D.; Panpan He, M.S.; Qinqin Li, M.S.; and Chengzhang Liu, M.S.

More information: *Hypertension* (2022).

www.ahajournals.org/doi/10.116 ... TENSIONAHA.121.18222

Provided by American Heart Association

Citation: Eating protein from a greater variety of sources may lower risk of high blood pressure (2022, March 10) retrieved 13 February 2024 from

<https://medicalxpress.com/news/2022-03-protein-greater-variety-sources-high.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|