


Gum disease is linked with heart attacks and diabetes

June 20 2022



Credit: Pixabay/CC0 Public Domain

A large study has found that people with a prior heart attack or diabetes are more likely to have gum disease than their healthy counterparts. The research is presented at [EuroPerio10](#), the world's leading congress in

periodontology and implant dentistry organized by the European Federation of Periodontology (EFP).

"This was an [observational study](#) and does not imply causal relationships," said study author Dr. Ida Stødle of the University of Oslo, Norway. "However, the findings raise awareness about the correlations between [chronic illnesses](#) which affect large numbers of people. This knowledge may help efforts to prevent these diseases."

Non-[communicable diseases](#) are becoming increasingly common as populations age and there is growing evidence that they are connected. Gum disease, also called periodontitis, affects up to half of all adults worldwide. This study examined whether severe periodontitis was more frequent in individuals with [diabetes](#) or a prior heart attack.

The analysis included 4,933 randomly selected participants of the community-based Trøndelag Health Study (HUNT Study). Participants completed questionnaires on sociodemographic and lifestyle factors, medications, and diseases such as type 2 diabetes and myocardial infarction. A clinical assessment of teeth and [soft tissues](#) was performed, plus a dental radiological examination. Weight, height, [blood pressure](#), and serum levels of cholesterol and glycated hemoglobin (HbA1c) were measured.

Dr. Stødle explained: "The presence of diabetes was assessed from self-reported questionnaires and may include a broad spectrum of severity, from poorly controlled to well controlled. For this reason we also examined the relationship between gum disease and HbA1c, which indicates average blood sugar levels over the last two to three months. Patients with diabetes are at higher risk of diabetic complications when their HbA1c levels are 48mmol/mol or above."

The average age of participants was 52 years and 56% were women. A

total of 147 (3.0%) participants reported a prior heart attack, 224 (4.5%) stated they had diabetes, 165 (3.3%) had elevated HbA1c (48mmol/mol or above), and 866 (17.6%) had severe periodontitis.

The researchers analyzed whether diabetes, elevated HbA1c, and prior heart attack predicted the likelihood of having severe gum disease. The analyses were adjusted for age, smoking, serum cholesterol, waist circumference and physical activity. The investigators found significant associations between diabetes, elevated HbA1c, prior heart attack and severe gum disease with odds ratios of 1.4, 1.5 and 1.7, respectively.

Dr. Stødle said: "The results show that patients with diabetes were 40% more likely to have severe gum disease than those without diabetes. Participants with high HbA1c were 50% more likely to have severe gum disease than those with HbA1c levels below 48mmol/mol. Finally, heart attack survivors were 70% more likely to have severe gum disease than participants who had never experienced a heart attack."

She concluded: "This study builds on previous evidence suggesting that people with gum disease are at greater risk of having a heart attack and developing diabetes, and also that those with diabetes are at greater risk of getting [gum disease](#). Taken together, the findings indicate that maintaining [oral health](#) could also benefit general health."

More information: HbA1c-level, diabetes and myocardial infarction are associated with periodontitis stage III and stage IV; The HUNT Study, presented 17 June at EuroPerio10: www.efp.org/europerio/

Provided by European Federation of Periodontology (EFP)

Citation: Gum disease is linked with heart attacks and diabetes (2022, June 20) retrieved 20

January 2023 from <https://medicalxpress.com/news/2022-06-gum-disease-linked-heart-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.