

Inability to stand on one leg for 10 seconds in mid to later life linked to near doubling in risk of death

June 21 2022



Credit: Pixabay/CC0 Public Domain

The inability to stand on one leg for 10 seconds in mid- to later life is linked to a near doubling in the risk of death from any cause within the next 10 years, finds research published online in the *British Journal of Sports Medicine*.

This simple and safe [balance](#) test could be included in routine health checks for older adults, say the researchers.

Unlike aerobic fitness and muscle strength and flexibility, balance tends to be reasonably well preserved until the sixth decade of life, when it starts to wane relatively rapidly, note the researchers.

Yet balance assessment isn't routinely included in health checks of [middle-aged](#) and older men and women, possibly because there isn't any standardized test for it, and there are few hard data linking it to clinical outcomes other than falls, they add.

The researchers therefore wanted to find out whether a balance test might be a reliable indicator of a person's risk of death from any cause within the next decade, and, as such, might therefore merit inclusion in routine health checks in later life.

The researchers drew on participants in the CLINIMEX Exercise cohort study. This was set up in 1994 to assess associations between various measures of [physical fitness](#), exercise-related variables, and conventional cardiovascular risk factors, with ill health and death.

The current analysis included 1,702 participants aged 51-75 (average of 61) at their first checkup, between February 2009 and December 2020. Around two thirds (68%) were men.

Weight and several measures of skinfold thickness plus waist size were taken. Details of medical history were also provided. Only those with stable gait were included.

As part of the checkup, participants were asked to stand on one leg for 10 seconds without any additional support.

To improve standardization of the test, participants were asked to place the front of the free foot on the back of the opposite lower leg, while keeping their arms by their sides and their gaze fixed straight ahead. Up to three attempts on either foot were permitted.

In all, around 1 in 5 (20.5%; 348) participants failed to pass the test. The inability to do so rose in tandem with age, more or less doubling at subsequent 5 year intervals from the age of 51-55 onwards.

The proportions of those unable to stand on one leg for 10 seconds were: nearly 5% among 51-55 year-olds; 8% among 56-60 year-olds; just under 18% among 61-65 year-olds; and just under 37% among 66-70 year-olds.

More than half (around 54%) of those aged 71-75 were unable to complete the test. In other words, people in this age group were more than 11 times as likely to fail the test as those just 20 years younger.

During an average monitoring period of 7 years, 123 (7%) people died: cancer (32%); cardiovascular disease (30%); respiratory disease (9%); and COVID-19 complications (7%).

There were no clear temporal trends in the deaths, or differences in the causes, between those able to complete the test and those who weren't able to do so.

But the proportion of deaths among those who failed the test was significantly higher: 17.5% vs. 4.5%, reflecting an absolute difference of just under 13%.

In general, those who failed the test had poorer health: a higher proportion were obese, and/or had heart disease, high blood pressure, and unhealthy blood fat profiles. And type 2 diabetes was 3 times as

common in this group: 38% vs. around 13%.

After accounting for age, sex, and underlying conditions, an inability to stand unsupported on one leg for 10 seconds was associated with an 84% heightened risk of death from any cause within the next decade.

This is an observational study, and as such, can't establish cause. As participants were all white Brazilians, the findings might not be more widely applicable to other ethnicities and nations, caution the researchers.

And information on potentially influential factors, including recent history of falls, physical activity levels, diet, smoking and the use of drugs that might interfere with balance wasn't available.

Nevertheless, the researchers conclude that the 10-second [balance test](#) "provides rapid and objective feedback for the patient and health professionals regarding static balance," and that the test "adds useful information regarding [mortality risk](#) in middle-aged and older men and women."

More information: Claudio Gil Araujo et al, Successful 10-second one-legged stance performance predicts survival in middle-aged and older individuals, *British Journal of Sports Medicine* (2022). [DOI: 10.1136/bjsports-2021-105360](#)

Provided by British Medical Journal

Citation: Inability to stand on one leg for 10 seconds in mid to later life linked to near doubling in risk of death (2022, June 21) retrieved 16 February 2023 from <https://medicalxpress.com/news/2022-06-inability-leg-seconds-mid-life.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.