


# Oncology therapy at the end-of-life: Have we missed the mark?

October 21 2022



Credit: *JAMA Oncology* (2022). DOI: 10.1001/jamaoncol.2022.4666

Continued treatment for patients with cancer near end-of-life delays conversations about goals of care and hospice enrollment, escalates costs, and may adversely affect the quality of care patients receive. Accordingly, over the last decade, major professional societies have recommended that clinicians decrease the use of systemic anti-cancer therapies at the end-of-life stage.

In a study published today in *JAMA Oncology*, researchers at Yale Cancer Center in collaboration with researchers from Flatiron Health, Inc., revealed that despite these recommendations, aggressive [cancer](#)

care at the end-of-life persists and there has been a substantial transition from the use of chemotherapy to immunotherapy.

"Over the last decade, the landscape of systemic anti-cancer [therapy](#) has changed dramatically following the approvals of multiple new targeted therapies," said Kerin Adelson, MD, Associate Professor of Medicine (Medical Oncology), Chief Quality Officer and Deputy Chief Medical Officer for Smilow Cancer Hospital, and senior author on the study.

"The quality measure that was developed ten years ago by professional oncology organizations, focuses on reducing end-of-life chemotherapy with the goal of providing patients with earlier palliative care. It is unclear whether this has benefited the end-of-life care patients receive."

To learn more, the researchers used the nationwide Flatiron Health electronic health record (EHR)-derived database to evaluate [adult patients](#) with cancer who received treatment and died within four years of diagnosis. Over the study period, rates of treatment within 30 days of death among all cancer types combined did not change (39% in 2015 and 2019) with similar trends seen for treatment within 14 days of death (17% in 2015 and 2019).

However, the type of systemic therapy received did vary; there were overall decreases in the use of chemotherapy (26% in 2015 and 16% in 2019) and increases in the use of immunotherapy (5% in 2015 and 18% in 2019). These changes were most noticeable in advanced non-small cell lung cancer and urothelial cancers where increases in treatment at end-of-life were driven by increased checkpoint inhibitor use.

"Our analysis identified no difference in the overall use of systemic anti-cancer therapy at the end-of-life since 2015. The approval of multiple new immunotherapy agents has engendered a great replacement phenomenon, substituting immunotherapy for chemotherapy," said Dr.

Adelson. "Increases in the use of targeted therapies may have interfered with our ability to achieve earlier palliative care integration and reduction in acute care utilization. More research is needed to determine if this change has affected the way [end-of-life](#) care is provided."

**More information:** Maureen E. Canavan et al, Systemic Anticancer Therapy at the End of Life—Changes in Usage Pattern in the Immunotherapy Era, *JAMA Oncology* (2022). [DOI: 10.1001/jamaoncol.2022.4666](#)

Provided by Yale Cancer Center

Citation: Oncology therapy at the end-of-life: Have we missed the mark? (2022, October 21) retrieved 28 February 2023 from <https://medicalxpress.com/news/2022-10-oncology-therapy-end-of-life.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.