

Statins stop hepatitis C virus replication

6 July 2006

Japanese scientists say they've found statins, typically used as anti-cholesterol medications, can inhibit the replication of the hepatitis C virus.

The findings mean statins might be able to replace ribavirin in combination therapy with interferon. There are 170 million people worldwide infected with HCV.

The standard HCV treatment is a combination therapy of interferon and ribavirin, which is effective in about 55 percent of patients. The remaining 45 percent face a threat of the disease progressing to cirrhosis and liver cancer.

Aware of recent studies showing one statin, lovastatin, inhibits HCV replication, researchers led by Masanori Ikeda of Okayama University tested other statins in search of a more effective anti-HCV therapy.

They evaluated the anti-HCV activities of five statins: atorvastatin, fluvastatin, lovastatin, pravastatin and simvastatin. When the statins were tested alone, all except pravastatin inhibited HCV replication, with fluvastatin having the strongest effect; atorvastatin and simvastatin had moderate effects and lovastatin had a weak effect.

The findings are reported in the July issue of the journal *Hepatology*.

Copyright 2006 by United Press International

APA citation: Statins stop hepatitis C virus replication (2006, July 6) retrieved 23 May 2021 from <https://medicalxpress.com/news/2006-07-statins-hepatitis-virus-replication.html>

<p><i>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</i></p>
