

# Folic acid cuts risk of cleft lip

January 26 2007

---

Taking folic acid supplements in early pregnancy seems to substantially reduce the risk of cleft lip, finds a new study published in the *British Medical Journal*.

It is widely known that taking folic acid in early pregnancy reduces the risk of neural tube defects, such as spina bifida. The current recommended daily dose is 400µg starting before conception.

Some studies have suggested that folic acid may also help prevent facial clefts, but the question remains unresolved. So researchers set out to assess the possible effects of folic acid on facial clefts in Norway, which has one of the highest rates of facial clefts in Europe.

They identified infants born from 1996 to 2000; 377 with cleft lip (with or without cleft palate), 196 with cleft palate only, and 763 healthy controls.

All mothers were surveyed about their reproductive history, smoking, alcohol, drugs, and other exposures during early pregnancy. They were also asked to recall their diet during the first three months of pregnancy, whether they took folic acid supplements and, if so, when and how often they took them.

Women were asked similar questions about multivitamins, and the researchers then estimated each woman's total folic acid intake.

After adjusting for smoking and other confounding factors, they found

that folic acid supplementation of 400µg or more a day reduced the risk of cleft lip with or without cleft palate by 40%.

Independent of supplements, diets rich in fruits, vegetables, and other high folate containing foods reduced the risk by 25%. The lowest risk of cleft lip was among women with folate rich diets who also took folic acid supplements and multivitamins. Folic acid provided no protection against cleft palate alone, according to the results.

The authors acknowledge that their study alone cannot show that folic acid definitely prevents cleft lip. Combined with all the previous evidence, however, their work does suggest a real preventive effect.

If folic acid is able to prevent a major birth defect in addition to neural tube defects, this benefit should be included among the risks and benefits of fortifying foods with folic acid, a matter of ongoing controversy in many countries, they conclude.

Source: British Medical Journal

Citation: Folic acid cuts risk of cleft lip (2007, January 26) retrieved 15 July 2023 from <https://medicalxpress.com/news/2007-01-folic-acid-cleft-lip.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--